

Antiferromagnetic gap in the Hubbard model

Tobias Baier, Eike Bick, and Christof Wetterich

Institut für Theoretische Physik

Universität Heidelberg

Philosophenweg 16, D-69120 Heidelberg

We compute the temperature dependence of the antiferromagnetic order parameter and the gap in the two dimensional Hubbard model at and close to half filling. Our approach is based on truncations of an exact functional renormalization group equation. The explicit use of composite bosonic degrees of freedom permits a direct investigation of the ordered low temperature phase. We show that the Mermin–Wagner theorem is not practically applicable for the spontaneous breaking of the continuous spin symmetry in the antiferromagnetic state. The critical behavior is dominated by the fluctuations of composite Goldstone bosons.

PACS numbers: 64.60.AK, 75.10.-b, 71.10.Fd, 74.20.-z

The low temperature phase of the Hubbard model [1] is highly interesting for a variety of phenomena observed in strongly correlated electron systems. This includes the antiferromagnetic state for zero or small doping, on which we concentrate in this note, as well as possible superconducting states for large doping. There has been a long debate about the effective theory for the spin wave fluctuations [2], [3] and the correct implementation of mean field theory [4, 5, 6], the problem being to reconcile the $SO(3)$ -spin symmetry with the Hartree-Fock results. More recently, functional renormalization group studies have made progress in the understanding of instabilities in various channels and the corresponding phase diagram [7, 8, 9, 10], but have been unable to explore the ordered phases for low temperature. (For other approaches see [11, 12].) We concentrate in this note on the two dimensional Hubbard model with next neighbor interactions. There a further problem arises for the understanding of the antiferromagnetic state - a theorem by Mermin and Wagner [13] forbids the spontaneous breaking of a continuous symmetry at nonvanishing temperature. This seems to contradict an antiferromagnetic order since the latter breaks spontaneously the continuous spin symmetry.

In this letter we present a combined study of Schwinger-Dyson (or Hartree-Fock) equations [14, 15], extended mean field methods [16], [17] using a generalized Hubbard-Stratonovich transformation [18] and functional renormalization based on an exact renormalization group equation [19, 20] for the effective average action [21]. We propose a partial bosonization which respects all symmetries. For a suitable choice the mean field theory (where only the fermionic fluctuations are taken into account) reproduces the Hartree-Fock result. For generic choices, however, the mean field phase diagram depends on unphysical parameters [16], reflecting the “Fierz ambiguity” [22] in the formulation of mean field theory. This problem is overcome by the inclusion of the bosonic fluctuations. The size of the bosonic corrections depends in such a way on the choice of partial bosonization that the phase diagram becomes almost independent of the choice of the mean field already for the approximations employed in the present work [23].

The scale dependence of the effective action is followed by the solution of a functional flow equation. This is formulated in the partially bosonized theory, employing the “rebosonization” technique of [24]. We thus combine the capability of mean field theory to describe an order parameter with the advantages of the functional renormalization group. The renormalization flow includes step by step the bosonic fluctuations on larger and larger length scales and the fermionic fluctuations closer and closer to the Fermi surface as the “infrared cutoff” k is lowered. This formulation of the functional renormalization group can be used in the low temperature phase as well, in contrast to the purely fermionic formulation [7, 8, 9, 10].

A coherent picture for the antiferromagnetic phase at low temperature and small doping emerges. We find a simple effective action for the spin waves (cf. eq. (15) and a more detailed discussion in [23]) which respects all symmetries. For temperatures near and below the effective critical temperature T_c and for small doping (small chemical potential μ) it can be described by an $O(3)$ -symmetric effective potential $U(\vec{a})$ for the antiferromagnetic order parameter $\vec{a}$ and an appropriate kinetic term which reads for small momenta $(Z_a t^2/2) \partial_i \vec{a} \partial_i \vec{a}$. (Here Z_a is a wave function renormalization, t is the next neighbor coupling of the Hubbard model and the derivatives ∂_i are in lattice units.) For $T < T_c$ the minimum of U occurs for $\alpha_0 = \vec{a}_0^2/2 \neq 0$. The characteristic scale for the antiferromagnetic order is set by the renormalized order parameter $\kappa_a = Z_a t^2 \alpha_0 / T$. In particular, this determines the gap for the electrons, $\Delta_a = \sqrt{2T\kappa_a/Z_a} (\bar{h}_a/t)$, with Yukawa coupling $\bar{h}_a/t$ close to one for our numerical example [23]. Also the renormalized mass (or inverse correlation length) for the spin fluctuations in the “radial direction” (parallel to the order parameter $\vec{a}_0$) is determined by κ_a , $\tilde{m}_a \sim \sqrt{\kappa_a}$, whereas the spin fluctuations in the “Goldstone direction” perpendicular to $\vec{a}_0$ are massless (infinite correlation length) for all $T < T_c$. Our result for the order parameter κ_a is plotted in fig. 1.

Actually, as a particularity of the two dimensional model, the size of the gap Δ_a (and the value of κ_a) depends on the infrared scale k which may be associated




FIG. 1: Renormalized expectation value κ_a of the antiferromagnetic condensate in the low temperature phase for $\mu = 0, U/t = 3$.

to the inverse of the size of the experimental probe. (We take $k^{-1} \approx 1\text{cm}$.) This is due to the fluctuations of the massless Goldstone bosons which reduce Δ_a as k is lowered. As we can see in fig. 2 this decrease is only logarithmic. For any fixed $T \neq 0$ and μ one always encounters a scale $k > 0$ where κ_a vanishes. Therefore, the order parameter vanishes for $k \rightarrow 0$, in agreement with the Mermin-Wagner theorem. Nevertheless, for low enough T the effective order parameter remains nonzero for any realistic macroscopic length k^{-1} . In this sense the Mermin-Wagner theorem fails to be practically applicable to the low temperature behavior of the two dimensional Hubbard model. For fixed macroscopic k^{-1} and $T < T_c$ the latter is well described by an ordered antiferromagnetic phase.

The effective critical temperature T_c depends mildly on the size of the probe k^{-1} - a change of k by a factor ten shifts T_c by 10%. Nevertheless, the two dimensional model exhibits effective critical behavior near T_c , with a macroscopic correlation length. For $k \ll T$ the fermion fluctuations effectively decouple and the flow is described by a classical $O(3)$ linear scalar model [25] (right part of fig. 2). For the correlation length ξ we find for $T > T_c$

$$\xi(T) = \frac{c(T)}{t} \exp \left\{ 20.7 \beta(T) \frac{T_c}{T} \right\} \quad (1)$$

with $T_c = 0.114t$, $\beta(T_c) = 1$ and $c(T), \beta(T)$ smoothly varying functions of T [23] without particular features for $T \rightarrow 0$. This closely resembles the result of [26] which would correspond to constant β . Near T_c the correlation length reaches the macroscopic size of the probe. Eq. (1) is no longer applicable for $T < T_c$.

The antiferromagnetic order occurs in the low temperature phase for a whole range of small doping. The critical temperature decreases mildly with the chemical potential μ [23]. We emphasize that the quantitative impact of the Goldstone-boson fluctuations on the value of T_c is very substantial. One may define a “pseudocritical tempera-




FIG. 2: Flow of the renormalized minimum of the potential κ_a (solid), the quartic bosonic coupling $10^{-2}\lambda_a$ (long dashes) and the wave function renormalization $10^{-1}Z_a$ (short dashes). We have chosen $U/t = 3$ and $T/t = 0.15$; $\mu = 0$.

ture” T_{pc} where the mass term for the antiferromagnetic spin wave reaches zero for some k , corresponding in the fermionic language to the four fermion coupling in the associated momentum channel growing “infinitely” large. We find $T_{pc}/t = 0.18(0.21)$ for $\mu = 0, T_{pc}/t = 0.145(0.18)$ for $\mu/t = 0.15$ (with brackets for mean field theory) - to be confronted with $T_c = 0.115$ for $\mu = 0$.

We next describe our formalism and approximations - for more details see [23]. We work within a functional integral formalism where the coordinates X denote the lattice sites as well as a “Euclidean time” τ . We concentrate in this note on fermion bilinears corresponding to spin density waves $\tilde{m}$ and charge density waves $\tilde{\rho}$.

$$\tilde{m}(X) = \hat{\psi}^\dagger(X) \vec{\sigma} \hat{\psi}(X), \quad \tilde{\rho}(X) = \hat{\psi}^\dagger(X) \hat{\psi}(X). \quad (2)$$

With these bilinears the four fermion coupling of the Hubbard model can be written as

$$(\hat{\psi}^\dagger(X) \hat{\psi}(X))^2 = \tilde{\rho}(X)^2 = -\frac{1}{3} \tilde{m}(X)^2. \quad (3)$$

We define the partially bosonized partition function by

$$Z[\eta, \eta^*, J_\rho, \vec{J}_m] = \int \mathcal{D}(\psi^*, \psi, \hat{\rho}, \hat{m}) \exp(- (S + S_\eta + S_J)) \quad (4)$$

with bosonic fields $\hat{m}, \hat{\rho}$, sources η, J and classical action

$$S = S_{F,\text{kin}} + \frac{1}{2} U_\rho \hat{\rho}^2 + \frac{1}{2} U_m \hat{m}^2 - U_\rho \hat{\rho} \tilde{\rho} - U_m \hat{m} \tilde{m}, \quad (5)$$

$$S_\eta = -\eta^\dagger \psi - \eta^T \psi^*, \quad S_J = -J_\rho \hat{\rho} - \vec{J}_m \hat{m}.$$

In particular, the fermion kinetic term in momentum space takes the form ($\omega_F = (2n+1)\pi T$)

$$S_{F,\text{kin}} = \sum_Q \hat{\psi}^\dagger(Q) P_F(Q) \hat{\psi}(Q), \quad (6)$$

$$P_F(Q) = i\omega_F + \epsilon - \mu, \quad \epsilon(\mathbf{q}) = -2t(\cos q_x + \cos q_y)$$

such that the Fermi surface is located at $-2t(\cos q_1 + \cos q_2) = \mu$. Here the Fourier transforms are defined as

$\hat{\psi}(X) = \sum_Q \hat{\psi}(Q) e^{iQX}$ and we use the short hand notation $Q = (\omega_F, \mathbf{q})$, $X = (\tau, \mathbf{x})$, $QX = \omega_F \tau + \mathbf{q} \cdot \mathbf{x}$.

The Gaussian functional integral over the spin density boson $\vec{m}(x)$ and charge density boson $\hat{\rho}(x)$ can be performed explicitly. This demonstrates that $W = \ln Z$ is equivalent to the free energy of the Hubbard model with next neighbor coupling and repulsive interaction $U > 0$ (up to terms $\sim J^2$ which are explicitly calculable). The equivalence holds for all U_ρ and U_m subject to the constraint

$$U = -U_\rho + 3U_m. \quad (7)$$

The remaining freedom in the choice of U_m is related to the Fierz ambiguity. The Hartree-Fock or lowest order Schwinger Dyson approximation corresponds to a computation of the fermionic fluctuation determinant (neglecting the fluctuations of $\hat{\rho}$, $\vec{m}$) for the choice $U_\rho = U_m = U/2$. (See [16] for a generalized setting).

The renormalization flow is implemented by the introduction of an infrared cutoff k through a piece in the action quadratic in the fields $\hat{\chi} = (\psi, \hat{\rho}, \vec{m}) = (\psi, \hat{b})$. In presence of this cutoff the generating functional for the connected Green functions depends on the scale k

$$W_k[J] = \ln \int \mathcal{D}\hat{\chi} \exp \left(- (S[\hat{\chi}] + \Delta S_k[\hat{\chi}] + J\hat{\chi}) \right), \quad (8)$$

$$\Delta S_k = \sum_Q \left\{ \hat{\psi}^\dagger(Q) R_k^\psi(Q) \psi(Q) + \frac{1}{2} \hat{b}(-Q) R_k^b(Q) \hat{b}(Q) \right\}.$$

For the fermionic cutoff we choose

$$R_k^\psi(Q) = i\omega_F \left(\frac{T_k}{T} - 1 \right) = 2\pi i (n_F + \frac{1}{2}) (T_k - T), \quad (9)$$

which has the effect of replacing the temperature T by some function T_k in the fermionic propagator. We specify this function by $T_k^4 = T^4 + k^4$, such that for large k the flow mimics a change in temperature whereas for $k \ll T$ the modification of the fermion propagator becomes ineffective. For the bosonic cutoff we take

$$R_k^b(Q) = Z(k^2 - \hat{Q}^2) \Theta(k^2 - \hat{Q}^2), \quad (10)$$

where $\hat{Q}^2 = Q^2$ in an appropriate interval, see eq. (16).

We define the *effective average action* as ($\chi = \partial W / \partial J$)

$$\Gamma_k[\chi] = J\chi - W_k[J] - \Delta S_k[\chi]. \quad (11)$$

The dependence on the scale k obeys an exact flow equation [19]

$$\partial_k \Gamma_k[\chi] = \frac{1}{2} \text{STr} \{ \partial_k R_k [\Gamma_k^{(2)} + R_k]^{-1} \}, \quad (12)$$

which has a physically intuitive “one loop form” [21]. Here the “supertrace” runs over field type, momentum and internal indices and has an additional minus sign for fermionic entries. We see that the IR cutoff R_k adds to

the exact inverse propagator matrix $\Gamma_k^{(2)}$ (second functional derivative) and therefore regulates possible zero modes, removing potential infrared problems for any $k > 0$. The effective average action interpolates between the classical action S (“initial condition” for $k \rightarrow \infty$) and the full “quantum effective action” for $k \rightarrow 0$.

As we are mainly interested in antiferromagnetic behavior we define the boson ($\Pi = (0, \pi, \pi)$)

$$\vec{a}(Q) = \vec{m}(Q + \Pi), \quad \vec{a}(X) = (-1)^P \psi^\dagger(X) \vec{\sigma} \psi(X), \quad (13)$$

with $P = 1$ for odd lattice sites and $P = 0$ else. The antiferromagnetic order corresponds to $\vec{a}(X)$ independent of X .

For an approximative solution of the functional differential equation (12) we employ a truncation of Γ_k consisting of a fermionic kinetic term, a Yukawa interaction between the fermions and bosons and a bosonic effective action for the $\vec{a}$ -boson. We describe here the choice $U_\rho = 0$, $U_m = U/3$ whereas an extended truncation including the fluctuations of the ρ -bosons can be found in [23]. For the fermionic kinetic term we adopt the classical part unchanged $\Gamma_{\psi,k} = S_{F,kin}(5)$. The Yukawa coupling $\bar{h}_{a,k}$ is taken to be scale dependent

$$\Gamma_{Y,k}[\psi, \psi^*, \vec{a}] = -\bar{h}_{a,k} \sum_{KQ} \vec{a}(K) \psi^*(Q) \vec{\sigma} \psi(Q') \quad (14)$$

$$\times \delta(K - Q + Q' + \Pi)$$

and the purely bosonic part is described by a kinetic term and an effective potential

$$\Gamma_{a,k}[\vec{a}] = \frac{1}{2} \sum_Q \vec{a}(-Q) P_a(Q) \vec{a}(Q) + \sum_X U_k(\alpha(X)). \quad (15)$$

Due to $SO(3)$ spin symmetry the k -dependent effective potential U_k only depends on the rotation invariant combination $\alpha(X) = \frac{1}{2} \vec{a}(X) \vec{a}(X)$. The boson kinetic term ($\omega_B = 2\pi mT$)

$$P_a(Q) = Z_{a,k} \hat{Q}^2 = Z_{a,k} (\omega_B^2 + t^2 [\mathbf{q}]^2), \quad (16)$$

involves a scale dependent wave function renormalization $Z_{a,k}$ (with the dimension of mass⁻¹) and the function $[\mathbf{q}]^2$ is defined as $[\mathbf{q}]^2 = q_x^2 + q_y^2$ for $q_i \in [-\pi, \pi]$ and continued periodically otherwise.

The flow equation for the effective potential $U_k(\alpha)$ obtains by evaluating eq. (12) for a homogeneous antiferromagnetic order parameter, $\vec{a}(Q) = \vec{a} \delta(Q)$, $\alpha = \frac{1}{2} \vec{a}^2$. The contribution of the fermionic fluctuations can be found by replacing in the mean field theory result $T \rightarrow T_k$ in the fermionic propagator and applying the formal derivative $\tilde{\partial}_k = (\partial T_k / \partial k) \partial / \partial T_k$. The bosonic contribution is the same as for the $O(3)$ linear σ -model [20, 21]:

$$\partial_k U(\alpha) = -2T \int_{-\pi}^{\pi} \frac{d^2 q}{(2\pi)^2} \tilde{\partial}_k \ln \cosh y(\alpha) \quad (17)$$

$$+ \frac{1}{2} \sum_{Q,i} \tilde{\partial}_k \ln [P_a(Q) + \hat{M}_i^2(\alpha) + R_k^a(Q)]$$

Here the squared fermion mass (gap) $\Delta_a^2 = 2\bar{h}_a^2\alpha$ enters the function

$$y(\alpha) = \frac{1}{2T_k} \sqrt{\epsilon^2(\mathbf{q}) + 2\bar{h}_a^2\alpha}. \quad (18)$$

The boson mass terms $\hat{M}_i^2$ obtain from the second derivatives of U_k and are different for the “radial mode” (in the direction of $\vec{a}$) and the “Goldstone modes” (perpendicular to $\vec{a}$). In the SSB regime we recognize two massless Goldstone bosons for α at the minimum of the potential, $\alpha = \alpha_0 \neq 0$.

The flow of Z_a is determined by the anomalous dimension $\eta_a = -k\partial_k \ln Z_a$,

$$\eta_a = h_a^2 t k (\partial_k T_k) \left[\frac{\partial}{\partial T_k} \frac{\partial}{\partial (l^2)} \right] \left\{ \frac{1}{T_k} \int_{-\pi}^{\pi} \frac{d^2 q}{(2\pi)^2} \frac{\tanh \frac{\epsilon(\mathbf{q}) + \mu}{2T_k} + \tanh \frac{\epsilon(\mathbf{q} + l\hat{e}_1) - \mu}{2T_k}}{(\epsilon(\mathbf{q}) + \epsilon(\mathbf{q} + l\hat{e}_1))/t} \right\} \Big|_{l=0}. \quad (19)$$

For the running of the renormalized Yukawa coupling $h_a = (T/Z_a)^{1/2} t^{-2} \bar{h}_a$ we obtain in the symmetric regime ($\alpha_0 = 0$) a direct and a “re-bosonized” contribution

$$\begin{aligned} k\partial_k h_a^2 &= h_a^2 \eta_a + \beta_{h_a^2}^{(d)} + \beta_{h_a^2}^{(rb)}, \\ \beta_{h_a^2}^{(d)} &= -2h_a^4 \frac{t^2}{kT} \tilde{\partial}_k \sum_Q \{ p_a^{-1}(Q) p_F^{-1}(Q) p_F^{-1}(Q + \Pi) \} \\ \beta_{h_a^2}^{(rb)} &= 2m_a^2 h_a^4 \frac{t^2}{kT} \tilde{\partial}_k \sum_Q \{ p_a^{-1}(Q) p_a^{-1}(\Pi - Q) \\ &\quad p_F^{-1}(-Q) [p_F^{-1}(\Pi - Q) - p_F^{-1}(Q)] \} \end{aligned} \quad (20)$$

with $m_a^2 = \partial U_k / \partial \alpha|_{\alpha=0}$, $p_F(Q) = (i\hat{\omega}_F - \mu + \epsilon(\mathbf{q}))/t$, $\hat{\omega}_F = (2n+1)\pi T_k$, $p_a(Q) = Q_k^2/k^2 + m_a^2$, $Q_k^2 = \hat{Q}^2 \Theta(Q^2 - k^2) + k^2 \Theta(k^2 - \hat{Q}^2)$. In the SSB-regime ($\alpha_0 \neq 0$) the change of the Yukawa coupling is negligible as we have checked numerically. Our results obtain from a numerical solution of the flow equations (17), (19), (20) with a quartic approximation to $U_k(\vec{a})$ [23]. The initial conditions for very large k are given by $\bar{h}_a = U_m = U/3$, $Z_a = 0$, $U_k(\alpha) = U_m \alpha$.

We believe that our truncation catches the most important features in the region of small doping. Our method gives a unified description which covers at once several interesting limiting cases: perturbation theory and the perturbative renormalization flow of the four-fermion coupling (small U), Hartree-Fock ($Z_a = 0$) and the classical spin wave description ($k \ll T$). The universal long-distance features (e.g. eq. (1)) are quite robust and should apply to a large class of models, including realistic materials: for decreasing T the correlation length increases so rapidly that it reaches the macroscopic size of the probe k^{-1} at T_c . For $T < T_c$ the formal symmetry restoration would require the randomization of antiferromagnetic domains with size much larger than k^{-1} . This effect is absent in a real experiment and macroscopic order can therefore be observed.

The analysis can be extended to include other bosonic channels, including the superconducting channel relevant for large doping [16], [23]. This would also lead to a more complete treatment of the momentum dependence of the effective four fermion interaction.

-
- [1] J. Hubbard, Proc. Roy. Soc., **A276**, 238, (1963); J. Kanamori, Prog. Theor. Phys., **30**, 275, (1963); M. C. Gutzwiller, Phys. Rev. Lett., **10**, 159, (1963)
 - [2] B. I. Shraiman, E. D. Siggia, Phys. Rev. Lett. **61**, 467 (1988)
 - [3] R. Shankar, Phys. Rev. Lett. **63**, 203 (1989)
 - [4] D. R. Penn, Phys. Rev. **142**, 350 (1966)
 - [5] A. A. Gomes, P. Lederer, J. Phys. (Paris) **38**, 231 (1977)
 - [6] H. J. Schulz, Phys. Rev. Lett. **65**, 2462 (1990)
 - [7] D. Zanchi and H. J. Schulz, Z.Phys. **B103**, 339 (1997); Europhys. Lett. **44**, 235 (1998)
 - [8] C. J. Halboth, W. Metzner, Phys. Rev. **B61**, 7364, (2000)
 - [9] C. Honerkamp, M. Salmhofer, cond-mat/0105218; C. Honerkamp, M. Salmhofer, T.M. Rice, cond-mat/0204063
 - [10] I. Grote, E. Kording, F. Wegner, cond-mat/0106604
 - [11] P. van Dongen, Phys. Rev. Lett. **67**, 757 (1991); Phys. Rev. **B50**, 14016 (1994)
 - [12] D.J. Scalapino, Phys.Rep. **250**, 329 (1995)
 - [13] N.D. Mermin, H. Wagner, Phys. Rev. Lett. **17**, 1133, (1966)
 - [14] F.J. Dyson, Phys. Rev. **75**, 1736, (1949); J. Schwinger, Proc. Nat. Acad. Sc., **37**, 452, (1951)
 - [15] J.W. Negele, H. Orland, *Quantum Many-Particle Systems*, Addison Wesley, Redwood City, 1988
 - [16] T. Baier, E. Bick, C. Wetterich, Phys. Rev., **B62**, 23, 15471, (2000) [cond-mat/0005218]
 - [17] T. Baier, E. Bick and C. Wetterich, [cond-mat/0107390]
 - [18] J. Hubbard, Phys. Rev. Lett. **3**, 77, (1959) ; R.L. Stratonovich, Sov. Phys. Dokl. **2**, 416, (1958)
 - [19] C. Wetterich, Phys. Lett. **B301**, 90, (1993)
 - [20] J. Berges, N. Tetradis, C. Wetterich, Phys.Rept. **363**, 223 (2002) [hep-ph/0005122]
 - [21] C. Wetterich, Nucl. Phys., **B352**, 529, (1991); Z. Phys., **C57**, 451, (1993); **C60**, 461, (1993)
 - [22] J. Jackel, C. Wetterich, [hep-ph/0207094], to appear in Phys. Rev. **D**
 - [23] T. Baier, E. Bick, C. Wetterich, cond-mat/0309715
 - [24] H. Gies, C. Wetterich, Phys.Rev. **D65**, 065001 (2002) [hep-th/0107221]
 - [25] G.v.Gersdorff, C. Wetterich, Phys. Rev. B **64**, 054513(2001) [hep-th/0008114].
 - [26] S. Chakravarty, B. Halperin, D. Nelson, Phys. Rev. **B39**, 2344 (1989)